



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

69. Proposed by GEORGE LILLEY, Ph. D., LL. D., Professor of Mathematics, University of Oregon, Eugene, Oregon.

An elliptic fence encloses a field whose major and minor axes are  $2a$  and  $2b$ , respectively. The ends of a rope, the length of which is equal to the length of the fence, are fastened outside the fence and at the extremities of the major axis. A horse is tethered by means of a ring which slides freely on the rope. Over how much ground can he feed? What is the length of the outside border? Find these values in square feet and feet, true to six decimal places, when the area of the field is one acre and  $a=2b$ .

Solution by MELLEN WOODMAN HASKELL, A. M., Ph. C., Associate Professor of Mathematics, University of California, Berkeley, Cal.

The outside border will evidently be a curve parallel to the given ellipse, so that the two curves will have common normals. Let  $ds$  denote the element of arc on the ellipse,  $dS$  the element of arc on the parallel curve lying between the same normals,  $d\phi$  the angle between those normals,  $dA$  the element of area bounded by those normals, by  $ds$  and  $dS$ , and  $p$  the length of the rope, then

$$dS=ds+pd\phi, \text{ and } dA=pd\phi+\frac{1}{2}p^2d\phi.$$

Integrating around the ellipse, since in this case the perimeter of the ellipse  $\int ds$  is equal to  $p$ , and also the complete integral  $\int d\phi$  is evidently  $2\pi$ , we have

$$S=p(1+2\pi) \text{ and } A=p^2(1+\pi).$$

A simple calculation then gives

$$\begin{aligned} p &= 806.693 \text{ feet, perimeter of ellipse;} \\ S &= 5875.293 \text{ feet, length of outside border;} \\ A &= 2695155 \text{ square feet, included area.} \end{aligned}$$

## MECHANICS.

57. Proposed by J. C. NAGLE, M. A., M. C. E., Professor of Civil Engineering, Agricultural and Mechanical College of Texas, College Station, Texas.

Over the intersection of two inclined planes slides a cord of uniform mass throughout its length. Find the equation to the path described by its center of gravity.

Solution by the PROPOSER.

Let  $AB, BC$  be the inclined planes,  $EBF$  the position of the cord at any instant. Let the length of the cord be  $a$ ; let  $BC$  be the  $x$ -axis,  $AB$  the  $y$ -axis. Let  $EB=z$ . Then the center of gravity of  $EB$  will be at  $G$ , the mid-point of  $EB$ , and of  $BF$  at  $M$ , the mid-point of  $BF$ . Let  $L$  be the center of gravity of whole cord. By moments about  $M$ ,

$$GM \times z = LM \times a, \text{ or } GM/LM = a/z \dots \dots \dots (1).$$

But from similar triangles,

$$\frac{GM}{LM} = \frac{BM}{HM} = \frac{\frac{1}{2}(a-z)}{\frac{1}{2}(a-z) - x} = a/z.$$

